

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 464 701 A1

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 158(3) EPC

(43) Date of publication:
06.10.2004 Bulletin 2004/41

(51) Int Cl.7: **C12N 15/09, C07K 14/435,
C07K 19/00, G01N 33/68**

(21) Application number: 02788859.3

(86) International application number:
PCT/JP2002/013363

(22) Date of filing: 20.12.2002

(87) International publication number:
WO 2003/054191 (03.07.2003 Gazette 2003/27)

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
IE IT LI LU MC NL PT SE SI SK TR**
Designated Extension States:
AL LT LV MK RO

(30) Priority: 20.12.2001 JP 2001387510

(71) Applicants:
• **Riken**
Wako-shi, Saitama 351-0198 (JP)
• **Medical & Biological Laboratories Co., Ltd.**
Nagayo-shi, Aichi 460-0002 (JP)

(72) Inventors:
• **MIYAWAKI, Atsushi**
Wako-shi, Saitama 351-0198 (JP)
• **KARASAWA, S.**
Medical & Biolg. Lab. Co. Ltd. Inalab
Ina-s hi, Nagano 396-0002 (JP)

(74) Representative:
Sternagel, Fleischer, Godemeyer & Partner
Patentanwälte
An den Gärten 7
51491 Overath (DE)

(54) FLUORESCENT PROTEINS

(57) An object of the present invention is to provide a novel fluorescent protein derived from organisms other than *Aequorea victoria*. According to the present invention, there is provided a fluorescent protein derived from *Fungia sp.*, which has the following properties:

- (1) the excitation maximum wavelength is 455 nm, and the fluorescence maximum wavelength is 488 nm;
- (2) the molar absorption coefficient at 455 nm is 38700 or 27700;
- (3) the quantum yield is 0.85 or 0.81; and
- (4) the pH sensitivity of the fluorescent property is stable at pH 5 to 9; and a fluorescent protein derived from *Fungia sp.*, which has the following properties:

- (1) the excitation maximum wavelength is 548 nm, and the fluorescence

- maximum wavelength is 561 nm;
- (2) the molar absorption coefficient at 548 nm is 75900 or 51000;
- (3) the quantum yield is 0.44 or 0.50; and
- (4) the pH sensitivity of the fluorescent property is pKa < 5.0.

EP 1 464 701 A1

Description

TECHNICAL FIELD

5 [0001] The present invention relates to a novel fluorescent protein having improved properties. More specifically, the present invention relates to a novel fluorescent protein derived from *Fungia sp.*, and the use thereof.

BACKGROUND ART

10 [0002] Green fluorescent protein (GFP) derived from *Aequorea victoria*, a jellyfish, has many purposes in biological systems. Recently, various GFP mutants have been produced based on the random mutagenesis and semi-rational mutagenesis, wherein a color is changed, a folding property is improved, luminance is enhanced, or pH sensitivity is modified. Fluorescent proteins such as GFP are fused with other proteins by gene recombinant technique, and monitoring of the expression and transportation of the fusion proteins is carried out.

15 [0003] One of the most commonly used types of GFP mutant is Yellow fluorescent protein (YFP). Among *Aequorea*-derived GFP mutants, YFP exhibits the fluorescence with the longest wavelength. The values ϵ and Φ of the majority of YFPs are 60,000 to 100,000 M⁻¹cm⁻¹ and 0.6 to 0.8, respectively (Tsien, R Y. (1998). Ann. Rev. Biochem. 67, 509-544). These values are comparable to those of the general fluorescent group (fluorescein, rhodamine, etc.). Accordingly, improvement of the absolute luminance of YFP is nearly approaching its limit.

20 [0004] Moreover, Cyan fluorescent protein (CFP) is another example of GFP mutants. Among such Cyan fluorescent proteins, ECFP (enhanced cyan fluorescent protein) has been known. Furthermore, Red fluorescent protein (RFP) has been isolated from sea anemone (*Discoma sp.*), and among such red fluorescent proteins, DasRed has been known. Thus, 4 types of fluorescent proteins including green, yellow, cyan and red fluorescent proteins, have been developed one after another, and their spectrum range has been significantly extended.

DISCLOSURE OF THE INVENTION

25 [0005] An object of the present invention is to provide a novel fluorescent protein derived from organisms other than *Aequorea victoria* and sea anemone. Another object of the present invention is to provide a novel fluorescent protein having improved fluorescent properties as compared with the fluorescent protein derived from *Aequorea victoria* or sea anemone.

30 [0006] In order to achieve the above objects, the present inventors have noticed fluorescent corals, and have conducted intensive studies to obtain a gene of fluorescent protein from corals by using suitable primers designed based on the amino acid sequences of known fluorescent proteins. As a result, they have succeeded in amplifying and cloning genes of fluorescent proteins from the cDNA library of *Fungia sp.*, a coral which is different from *Galaxea fascicularis* from which a gene of a fluorescent protein was previously obtained, using the above primers. Further, the present inventors have examined the fluorescent properties of the obtained fluorescent proteins derived from *Fungia sp.* and as a result, they have found that these fluorescent proteins have desired fluorescent properties. The present invention has been completed based on these findings.

35 [0007] Thus, the present invention provides a fluorescent protein derived from *Fungia sp.*, which has the following properties:

- 40 (1) the excitation maximum wavelength is 455 nm, and the fluorescence maximum wavelength is 488 nm;
 (2) the molar absorption coefficient at 455 nm is 38700 or 27700;
 45 (3) the quantum yield is 0.85 or 0.81; and
 (4) the pH sensitivity of the fluorescent property is stable at pH 5 to 9.

50 [0008] The present invention further provides a fluorescent protein derived from *Fungia sp.*, which has the following properties:

- (1) the excitation maximum wavelength is 548 nm, and the fluorescence maximum wavelength is 561 nm;
 (2) the molar absorption coefficient at 548 nm is 75900 or 51000;
 (3) the quantum yield is 0.44 or 0.50; and
 55 (4) the pH sensitivity of the fluorescent property is pKa < 5.0.

[0009] In another aspect of the present invention, there is provided a fluorescent protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2; or

(b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1 or 2, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2.

5 [0010] Further, in another aspect of the present invention, there is provided a fluorescent protein of the following (a) or (b):

(a) a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4; or
 (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several
 10 amino acids with respect to the amino acid sequence shown in SEQ ID NO: 3 or 4, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4.

[0011] In further another aspect of the present invention, there is provided a protein having an amino acid sequence wherein cysteine at position 64 is substituted with alanine with respect to the amino acid sequence shown in SEQ ID
 15 NO: 3, or an amino acid sequence wherein glutamic acid at position 211 is substituted with alanine with respect to the amino acid sequence shown in SEQ ID NO: 3.

[0012] In further another aspect of the present invention, there is provided DNA which encodes the fluorescent protein of the present invention as mentioned above.

20 [0013] In further another aspect of the present invention, there is provided DNA which encodes the protein of the following (a) or (b):

(a) a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2; or
 (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several
 25 amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1 or 2, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2.

[0014] In further another aspect of the present invention, there is provided DNA which encodes the protein of (a) or (b):

(a) a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4; or
 30 (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 3 or 4, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4.

35 [0015] In further another aspect of the present invention, there is provided DNA of the following (a) or (b):

(a) DNA having a nucleotide sequence shown in SEQ ID NO: 5 or 6; or
 (b) DNA having a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucle-
 otides with respect of the nucleotide sequence shown in SEQ ID NO: 5 or 6, and encoding a protein having fluo-
 40 rescent properties which are equivalent to those of a protein encoded by the nucleotide sequence shown in SEQ ID NO: 5 or 6.

[0016] In further another aspect of the present invention, there is provided DNA of the following (a) or (b):

(a) DNA having a nucleotide sequence shown in SEQ ID NO: 7 or 8; or
 45 (b) DNA having a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucle-
 otides with respect of the nucleotide sequence shown in SEQ ID NO: 7 or 8, and encoding a protein having fluo-
 rescent properties which are equivalent to those of a protein encoded by the nucleotide sequence shown in SEQ ID NO: 7 or 8.

50 [0017] In further another aspect of the present invention, there is provided a recombinant vector having the above-described DNA of the present invention.

[0018] In further another aspect of the present invention, there is provided a transformant having the above-described DNA or recombinant vector of the present invention.

55 [0019] In further another aspect of the present invention, there is provided a fusion fluorescent protein consisting of the above-described fluorescent protein of the present invention and another protein. Preferably, said another protein is one that localizes in the cell, and more preferably, said another protein is one specific to an intracellular organella.

[0020] In further another aspect of the present invention, there is provided a method for analyzing the localization or dynamics of a protein in cells, characterized in that the above-described fusion protein of the present invention is

allowed to be expressed in cells.

[0021] In another aspect of the present invention, there is provided a fluorescent reagent kit which comprises the above-described fluorescent protein, DNA, recombinant vector, transformant or fusion protein according to the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022]

Figure 1 shows results obtained by comparing the fluorescence spectrum and excitation spectrum of fluorescent proteins derived from *Fungia sp.* of the present invention (KCy-1 and KCy-2) with those of ECFP, and by comparing the fluorescence spectrum and excitation spectrum of fluorescent proteins derived from *Fungia sp.* of the present invention (KO-1 and KO-2) and those of DsRed.

Figure 2 shows results obtained by analyzing the fluorescent properties of the fluorescent proteins KCy-1, KCy-2, KO-1 and KO-2 derived from *Fungia sp.* of the present invention.

(1) Difference in coloration between Kcy-1 and Kcy-2 in *Escherichia coli*
(Fluorescence at 488 nm obtained by excitation at 400 nm)

(2) Difference in coloration between KO-1 and KO-2 in *Escherichia coli*
(Fluorescence at 561 nm obtained by excitation at 500 nm)

(3) Change in fluorescence between KO-1 and KO-2
(The value (maturation value) of orange components (561 nm) / green components (508nm) obtained by excitation at 470 nm)

Figure 3 shows the change over time in the green components (508 nm) and orange components (561 nm) in fluorescent proteins (KO-1 and KO-2) derived from *Fungia sp.* of the present invention.

Figure 4 shows the pH sensitivity of the fluorescence intensity of each of the fluorescent proteins (KCy-1, KCy-2, KO-1 and KO-2) derived from *Fungia sp.* of the present invention.

Figure 5 shows the fluorescence spectrum (em) and excitation spectrum (ex) (left figure), and absorption spectrum (right figure) of a mutant (KO-C64A) of a fluorescent protein derived from *Fungia sp.* of the present invention.

Figure 6 shows the fluorescence spectrum (em) and excitation spectrum (ex) (left figure), and absorption spectrum (right figure) of a mutant (KO-E211A) of a fluorescent protein derived from *Fungia sp.* of the present invention.

BEST MODE FOR CARRYING OUT THE INVENTION

[0023] The embodiments of the present invention will be described in detail below.

(1) Fluorescent protein of the present invention

[0024] The fluorescent protein of the present invention is characterized in that it is derived from *Fungia sp.* and has the following properties:

- (1) the excitation maximum wavelength is 455 nm, and the fluorescence maximum wavelength is 488 nm;
- (2) the molar absorption coefficient at 455 nm is 38700 or 27700;
- (3) the quantum yield is 0.85 or 0.81; and
- (4) the pH sensitivity of the fluorescent property is stable at pH 5 to 9.

[0025] Another fluorescent protein of the present invention is characterized in that it is derived from *Fungia sp.* and has the following properties:

- (1) the excitation maximum wavelength is 548 nm, and the fluorescence maximum wavelength is 561 nm;
- (2) the molar absorption coefficient at 548 nm is 75900 or 51000;
- (3) the quantum yield is 0.44 or 0.50; and
- (4) the pH sensitivity of the fluorescent property is $pK_a < 5.0$.

[0026] *Fungia sp.* is one type of coral and mainly lives in the western region of the Atlantic Ocean. *Fungia sp.* is characterized in that the outer shape of a group of the corals is a polygon, having long tentacles, and that a group of the corals as a whole develops a clear orange color.

[0027] In Examples of the present specification described later, the fluorescent protein of the present invention having the above properties was isolated by using *Fungia sp.* as a starting material. However, in some cases, the fluorescent protein of the present invention can be obtained also from coral which emits fluorescence other than *Fungia sp.* Such fluorescent proteins are also included in the scope of the present invention.

5 [0028] As described in Examples mentioned later, the first fluorescent protein (KCy-1) of the present invention has an excitation maximum wavelength of 455 nm and a fluorescence maximum wavelength of 488 nm. It has a molar absorption coefficient of 38700 (455 nm) and a quantum yield of 0.85. As described in Examples mentioned later, the second fluorescent protein (KCy-2) of the present invention has an excitation maximum wavelength of 455 nm and a fluorescence maximum wavelength of 488 nm. It has a molar absorption coefficient of 27700 (455 nm) and a quantum yield of 0.81. In contrast, ECFP (Clontech) has a molar absorption coefficient of 28700 (435 nm) and a quantum yield of 0.40.

10 [0029] As described in Examples mentioned later, the third fluorescent protein (KO-1) of the present invention has an excitation maximum wavelength of 548 nm and a fluorescence maximum wavelength of 561 nm. It has a molar absorption coefficient of 75900 (548 nm) and a quantum yield of 0.44. As described in Examples mentioned later, the fourth fluorescent protein (KO-2) of the present invention has an excitation maximum wavelength of 548 nm and a fluorescence maximum wavelength of 561 nm. It has a molar absorption coefficient of 51000 (548 nm) and a quantum yield of 0.50. In contrast, DsRed (Clontech) has a molar absorption coefficient of 86100 (559 nm) and a quantum yield of 0.29.

15 [0030] Molar absorption coefficient represents the absorption amount of photons per mole of fluorescent molecules. Quantum yield is a numerical value showing the amount of the absorbed photons that can be emitted as fluorescence. Thus, when the values of the molar absorption coefficient and quantum yield are great, it shows that strong fluorescence is emitted. Accordingly, among the above-described fluorescent proteins of the present invention, since cyan fluorescent proteins KCy-1 and KCy-2 have a molar absorption coefficient and a quantum yield greater than those of ECFP, these proteins emit fluorescence stronger than that of ECFP. More specifically, KCy-1 emits fluorescence approximately 2 or 3 times brighter than that of ECFP, and KCy-2 emits fluorescence approximately 1.5 times brighter than that of ECFP.

20 [0031] With regard to the maximum excitation wavelength and the maximum fluorescence wavelength, there are no significant differences between ECFP and the fluorescent proteins KCy-1 and KCy-2 of the present invention. However, differing from ECFP, the excitation and fluorescence spectrums of the fluorescent proteins KCy-1 and KCy-2 of the present invention do not have a shoulder on their long wavelength side, but they have a sharp form. Accordingly, it can be said that these fluorescent proteins are advantageously used in multicolor imaging or the like which is performed in combination with other fluorescent molecules.

25 [0032] Moreover, KCy-1 and KCy-2 are characterized in that their fluorescence is low sensitive to pH in the range between pH 5 and pH 9. This is to say, there is only a slight extent of fluctuation in the peak value of their fluorescence intensity in the range between pH 5 and pH 9, and a high fluorescence intensity can be maintained in this pH range. In the case of the previously used ECFP, since its fluorescence intensity is decreased at pH 7 or lower, its use in living bodies has a certain limit. However, the fluorescent proteins of the present invention do not have such a limit.

30 [0033] On the other hand, orange fluorescent proteins KO-1 and KO-2 among the above-described fluorescent proteins of the present invention emit fluorescence approximately 2 times brighter than that of a red fluorescent protein (DsRed) derived from sea anemone (*Discosoma*). In addition, each of KO-1 and KO-2 has the peak of their fluorescence spectrum at a wavelength different from those of the existing fluorescent proteins. This is to say, EYFP (yellow) (Clontech) has the peak of its fluorescence spectrum around 530 nm, and DsRed (Clontech) has the peak of its fluorescence spectrum around 580 nm. In contrast, the KO-1 and KO-2 of the present invention has the peak of their fluorescence spectrum around 561 nm.

35 [0034] In addition, specific examples of a mutant of the protein KO-1 of the present invention may include:

- (1) a mutant which is obtained by substituting cysteine at position 64 by alanine in the amino acid sequence of KO-1, and emits green fluorescence (fluorescence maximum at 508 nm, and excitation maximum at 496 nm), wherein the fluorescent properties are sifted on the short wavelength side when compared with KO; and
- 40 (2) a mutant which is obtained by substituting glutamic acid at position 211 by alanine in the amino acid sequence of KO-1, and emits red fluorescence (fluorescence maximum at 578 nm, and excitation maximum at 563 nm), wherein fluorescent properties are sifted on the long wavelength side when compared with KO.

45 [0035] The examples of the fluorescent protein of the present invention include a fluorescent protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several

amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1 or 2, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2.

[0036] The further examples of the fluorescent protein of the present invention include a fluorescent protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 3 or 4, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4.

[0037] The scope of "one or several" in the phrase "an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids" used herein is not particularly limited in the present specification. For example, it means 1 to 20, preferably 1 to 10, more preferably 1 to 7, further preferably 1 to 5, and particularly preferably 1 to 3.

[0038] The term "fluorescent properties which are equivalent" used herein means that the protein has an equivalent fluorescence intensity, an equivalent excitation wavelength, an equivalent fluorescence wavelength, an equivalent pH sensitivity, or the like.

[0039] The method of obtaining the fluorescent protein of the present invention is not particularly limited. The protein may be either a protein synthesized by chemosynthesis, or recombinant protein produced by a gene recombination technique.

[0040] Where a recombinant protein is produced, it is necessary to obtain DNA encoding the protein. Appropriate primers are designed by utilizing information regarding the amino acid sequence shown in SEQ ID NOS: 1 to 4 of the sequence listing of the present specification and the nucleotide sequence shown in SEQ ID NOS: 5 to 8 thereof. Using these primers, PCR is carried out by using cDNA clones of the above-described various types of known fluorescent proteins as a template, so that DNA encoding the fluorescent protein of the present invention can be obtained. Where a partial fragment of DNA encoding the fluorescent protein of the present invention are obtained by the above-described PCR, the produced DNA fragments are ligated to one another by a gene recombination technique, so that DNA encoding the desired fluorescent protein can be obtained. The fluorescent protein of the present invention can be produced by introducing this DNA into an appropriate expression system. Expression in an expression system will be described later in the present specification.

(2) DNA of the present invention

[0041] According to the present invention, a gene encoding the fluorescent protein of the present invention is provided.

[0042] Specific examples of DNA encoding the fluorescent protein of the present invention may include DNA which encodes the protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1 or 2, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2.

[0043] Further specific examples of DNA encoding the fluorescent protein of the present invention may include DNA which encodes the protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 3 or 4, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4.

[0044] Further specific examples of DNA encoding the fluorescent protein of the present invention may include DNA of the following (a) or (b):

- (a) DNA having a nucleotide sequence shown in SEQ ID NO: 5 or 6; or
- (b) DNA having a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucleotides with respect of the nucleotide sequence shown in SEQ ID NO: 5 or 6, and encoding a protein having fluorescent properties which are equivalent to those of a protein encoded by the nucleotide sequence shown in SEQ

ID NO: 5 or 6.

[0045] Further specific examples of DNA encoding the fluorescent protein of the present invention may include DNA of the following (a) or (b):

- (a) DNA having a nucleotide sequence shown in SEQ ID NO: 7 or 8; or
 (b) DNA having a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucleotides with respect of the nucleotide sequence shown in SEQ ID NO: 7 or 8, and encoding a protein having fluorescent properties which are equivalent to those of a protein encoded by the nucleotide sequence shown in SEQ ID NO: 7 or 8.

[0046] The DNA of the present invention can be synthesized by, for example, the phosphoramidite method, or it can also be produced by polymerase chain reaction (PCR) using specific primers. The DNA of the present invention or a fragment thereof is produced by the method described above in the specification.

[0047] A method of introducing a desired mutation into a certain nucleic acid sequence is known to a person skilled in the art. For example, known techniques such as a site-directed mutagenesis, PCR using degenerated oligonucleotides, or the exposure of cells containing nucleic acid to mutagens or radioactive rays, are appropriately used, so as to construct DNA having a mutation. Such known techniques are described in, for example, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY., 1989; and Current Protocols in Molecular Biology, Supplements 1 to 38, John Wiley & Sons (1987-1997).

(3) Recombinant vector of the present invention

[0048] The DNA of the present invention can be inserted into a suitable vector and used. The type of a vector used in the present invention is not particularly limited. For example, it may be either a vector that can autonomously replicate (e.g., a plasmid, etc.), or vector that is incorporated into the genomes of host cells when it is introduced into the host cells and is then replicated together with the chromosome into which it is incorporated.

[0049] The vector used in the present invention is preferably an expression vector. In an expression vector, elements necessary for transcription (e.g., a promoter, etc.) are functionally ligated to the DNA of the present invention. The promoter is a DNA sequence which shows a transcriptional activity in host cells, and it is appropriately selected depending on the type of host cells.

[0050] Examples of a promoter which can operate in bacterial cells may include a *Bacillus stearothermophilus* maltogenic amylase gene promoter, a *Bacillus licheniformis* alpha-amylase gene promoter, a *Bacillus amyloliquefaciens* BAN amylase gene promoter, a *Bacillus subtilis* alkaline protease gene promoter, a *Bacillus pumilus* xylosidase gene promoter, P_R and P_L promoters of phage lambda, and lac, trp and tac promoters of *Escherichia coli*.

[0051] Examples of a promoter which can operate in mammalian cells may include an SV40 promoter, an MT-1 (metallothionein gene) promoter, and an adenovirus-2 major late promoter. Examples of a promoter which can operate in insect cells may include a polyhedrin promoter, a P10 promoter, an *Autographa californica* polyhedrosis basic protein promoter, a baculovirus immediate-early gene 1 promoter, and a baculovirus 39K delayed-early gene promoter. Examples of a promoter which can operate in yeast host cells may include promoters derived from yeast glycolytic genes, an alcohol dehydrogenase gene promoter, a TPI1 promoter, and an ADH2-4c promoter.

[0052] Examples of a promoter which can operate in filamentous cells may include an ADH3 promoter and a tpiA promoter.

[0053] In addition, an appropriate terminator such as a human growth hormone terminator, or a TPI1 terminator or ADH3 terminator for fungal cells, may be functionally bound to the DNA of the present invention, as necessary. The recombinant vector of the present invention may further have elements such as a polyadenylation signal (e.g., one derived from SV40 or the adenovirus 5E1b region), a transcription enhancer sequence (e.g., an SV40 enhancer), or a translation enhancer sequence (e.g., one encoding the adenovirus VA RNA).

[0054] The recombinant vector of the present invention may further comprise a DNA sequence which enables the replication of the recombinant vector in host cells. SV40 replication origin is an example of such a sequence (when the host cells are mammalian cells).

[0055] The recombinant vector of the present invention may further comprise a selective marker. Examples of such a selective marker may include genes, complements of which are absent from host cells, such as a dihydrofolate reductase (DHFR) gene or a *Shizosaccharomyces pombe* TPI gene, and drug resistant genes such as ampicillin, kanamycin, tetracycline, chloramphenicol, neomycin or hygromycin-resistant genes.

[0056] A method for ligating the DNA of the present invention, a promoter and, as desired, a terminator and/or a secretory signal sequence to one another and inserting these items into a suitable vector is known to a person skilled in the art.

(4) Transformant of the present invention

[0057] A transformant can be produced by introducing the DNA or recombinant vector of the present invention into a suitable host.

5 [0058] Any cell can be used as a host cell into which the DNA or recombinant vector of the present invention is introduced, as long as the DNA construct of the present invention can be expressed therein. Examples of such a cell may include bacteria, yeasts, fungal cells, and higher eukaryotic cells.

[0059] Examples of bacteria may include Gram-positive bacteria such as *Bacillus* or *Streptomyces*, and Gram-negative bacteria such as *Escherichia coli*. These bacteria may be transformed by the protoplast method or other known

10 [0060] Examples of mammalian cells may include HEK 293 cells, HeLa cells, COS cells, BHK cells, CHL cells, and CHO cells. A method of transforming mammalian cells and expressing the introduced DNA sequence in the cells is also known. Examples of such a method may include the electroporation, the calcium phosphate method, and the lipofection method.

15 [0061] Examples of yeast cells may include those belonging to *Saccharomyces* or *Shizosaccharomyces*. Examples of such cells may include *Saccharomyces cerevisiae* and *Saccharomyces kluyveri*. Examples of a method of introducing a recombinant vector into yeast host cells may include the electroporation, the spheroplast method, and the lithium acetate method.

20 [0062] Examples of other fungal cells may include those belonging to *Filamentous fungi* such as *Aspergillus*, *Neurospora*, *Fusarium* or *Trichoderma*. Where *Filamentous fungi* are used as host cells, transformation can be carried out by incorporating DNA constructs into host chromosomes, so as to obtain recombinant host cells. Incorporation of DNA constructs into the host chromosomes is carried out by known methods, and such known methods may include homologous recombination and heterologous recombination.

25 [0063] Where insect cells are used as host cells, both a vector into which a recombinant gene is introduced and a baculovirus are co-introduced into insect cells, and a recombinant virus is obtained in the culture supernatant of the insect cells. Thereafter, insect cells are infected with the recombinant virus, so as to allow the cells to express proteins (described in, for example, *Baculovirus Expression Vectors, A Laboratory Manual*; and *Current Protocols in Molecular Biology*, Bio/Technology, 6, 47 (1988)).

30 [0064] The *Autographa californica* nuclear polyhedrosis virus, which is a virus infecting to insects belonging to *Bathra brassicae*, can be used as baculovirus.

[0065] Examples of insect cells used herein may include Sf9 and Sf21, which are *Spodoptera frugiperda* ovarian cells [*Baculovirus Expression Vectors, A Laboratory Manual*, W. H. Freeman & Company, New York, (1992)], and HiFive (manufactured by Invitrogen), which are *Trichoplusia ni* ovarian cells.

35 [0066] Examples of the method of co-introducing both a vector into which a recombinant gene has been introduced and the above baculovirus into insect cells to prepare a recombinant virus, may include the calcium phosphate method and the lipofection method.

[0067] The above transformant is cultured in an appropriate nutritive medium under conditions enabling the introduced DNA construct to be expressed. In order to isolate and purify the fusion fluorescent protein of the present invention from the culture product of the transformant, common methods of isolating and purifying proteins may be used.

40 [0068] For example, where the protein of the present invention is expressed in a state dissolved in cells, after completion of the culture, cells are recovered by centrifugal separation, and the recovered cells are suspended in a water type buffer. Thereafter, the cells are disintegrated using an ultrasonic disintegrator or the like, so as to obtain a cell-free extract. A supernatant is obtained by centrifuging the cell-free extract, and then, a purified sample can be obtained from the supernatant by applying, singly or in combination, the following ordinary protein isolation and purification

45 methods: the solvent extraction, the salting-out method using ammonium sulfate or the like, the desalting method, the precipitation method using an organic solvent, the anion exchange chromatography using resins such as diethylaminoethyl (DEAE) sepharose, the cation exchange chromatography using resins such as S-Sepharose FF (manufactured by Pharmacia), the hydrophobic chromatography using resins such as butyl sepharose or phenyl sepharose, the gel filtration method using a molecular sieve, the affinity chromatography, the chromatofocusing method, and the electro-

50 phoresis such as isoelectric focusing.

(5) Use of the fluorescent protein of the present invention and a fusion fluorescent protein comprising the same

55 [0069] The fluorescent protein of the present invention can be fused with another protein, so as to construct a fusion fluorescent protein.

[0070] A method of obtaining the fusion fluorescent protein of the present invention is not particularly limited. It may be either a protein synthesized by chemosynthesis, or recombinant protein produced by a gene recombination technique.

[0071] Where a recombinant protein is produced, it is necessary to obtain DNA encoding the protein. Appropriate primers are designed by using the information regarding the amino acid sequence shown in SEQ ID NOS: 1 to 4 of the sequence listing of the present specification and the nucleotide sequence shown in SEQ ID NOS: 5 to 8 thereof. Using these primers, PCR is carried out using a DNA fragment containing the gene of the fluorescent protein of the present invention as a template, so as to produce DNA fragments necessary for construction of the DNA encoding the fluorescent protein of the present invention. Moreover, DNA fragments encoding a protein to be fused is also obtained in the same above manner.

[0072] Subsequently, the thus obtained DNA fragments are ligated to one another by a gene recombination technique, so that DNA encoding the desired fusion fluorescent protein can be obtained. This DNA is then introduced into an appropriate expression system, so that the fusion fluorescent protein of the present invention can be produced.

[0073] The fluorescent protein of the present invention has an extremely high utility value as a marker. This is to say, the fluorescent protein of the present invention is purified as a fusion protein with an amino acid sequence to be tested, and the fusion protein is introduced into cells by methods such as the microinjection. By observing the distribution of the fusion protein over time, targeting activity of the amino acid sequence to be tested can be detected in the cells.

[0074] The type of another protein (an amino acid sequence to be tested) with which the fluorescent protein of the present invention is fused is not particularly limited. Preferred examples may include proteins localizing in cells, proteins specific for intracellular organelles, and targeting signals (e.g., a nuclear transport signal, a mitochondrial presequence, etc.). In addition, the fluorescent protein of the present invention can be expressed in cells and used, as well as being introduced into cells by the microinjection or the like. In this case, a vector into which the DNA encoding the fluorescent protein of the present invention is inserted in such a way that it can be expressed, is introduced into host cells.

[0075] Moreover, the fluorescent protein of the present invention can also be used as a reporter protein to determine promoter activity. This is to say, a vector is constructed such that DNA encoding the fluorescent protein of the present invention is located downstream of a promoter to be tested, and the vector is then introduced into host cells. By detecting the fluorescence of the fluorescent protein of the present invention which is emitted from the cells, the activity of the promoter to be tested can be determined. The type of a promoter to be tested is not particularly limited, as long as it operates in host cells.

[0076] A vector used to detect the targeting activity of the above amino acid sequence to be tested or to determine promoter activity is not particularly limited. Examples of a vector preferably used for animal cells may include pNEO (P. Southern, and P. Berg (1982) J. Mol. Appl. Genet. 1: 327), pCAGGS (H. Niwa, K. Yamamura, and J. Miyazaki, Gene 108, 193-200 (1991)), pRc/CMV (manufactured by Invitrogen), and pCDM8 (manufactured by Invitrogen). Examples of a vector preferably used for yeasts may include pRS303, pRS304, pRS305, pRS306, pRS313, pRS314, pRS315, pRS316 (R S. Sikorski and P. Hieter (1989) Genetics 122: 19-27), pRS423, pRS424, pRS425, pRS426 (T. W. Christianson, R S. Sikorski, M. Dante, J. H. Shero, and P. Hieter (1992) Gene 110: 119-122).

[0077] In addition, the type of cells used herein is also not particularly limited. Various types of animal cells such as L cells, BalbC-3T3 cells, NIH3T3 cells, CHO (Chinese hamster ovary) cells, HeLa cells or NRK (normal rat kidney) cells, yeast cells such as *Saccharomyces cerevisiae*, *Escherichia coli* cells, or the like can be used. Vector can be introduced into host cells by common methods such as the calcium phosphate method or the electroporation.

[0078] The above obtained fusion fluorescent protein of the present invention wherein the fluorescent protein of the present invention is fused with another protein (referred to as a protein X) is allowed to be expressed in cells. By monitoring a fluorescence emitted, it becomes possible to analyze the localization or dynamics of the protein X in cells. That is, cells transformed or transfected with DNA encoding the fusion fluorescent protein of the present invention are observed with a fluorescence microscope, so that the localization and dynamics of the protein X in the cells can be visualized and thus analyzed.

[0079] For example, by using a protein specific for an intracellular organella as a protein X, the distribution and movement of a nucleus, a mitochondria, an endoplasmic reticulum, a Golgi body, a secretory vesicle, a peroxisome, etc., can be observed.

[0080] Moreover, for example, axis cylinders or dendrites of the nerve cells show an extremely complicated change in strikes in an individual who is under development. Accordingly, fluorescent labeling of these sites enables a dynamic analysis.

[0081] The fluorescence of the fluorescent protein of the present invention can be detected with a viable cell. Such detection can be carried out using, for example, a fluorescence microscope (Axiophoto Filter Set 09 manufactured by Carl Zeiss) or an image analyzer (Digital Image Analyzer manufactured by ATTO).

[0082] The type of a microscope can be appropriately selected depending on purposes. Where frequent observation such as pursuit of a change over time is carried out, an ordinary incident-light fluorescence microscope is preferable. Where observation is carried out while resolution is emphasized, for example, in the case of searching localization in cells specifically, a confocal laser scanning microscope is preferable. In terms of maintenance of the physiological state of cells and prevention from contamination, an inverted microscope is preferable as a microscope system. When an erecting microscope with a high-powered lens is used, a water immersion lens can be used.

[0083] A filter set can be appropriately selected depending on the fluorescence wavelength of a fluorescent protein. In the case of the fluorescent protein having the excitation maximum wavelength of 455 nm and the fluorescence maximum wavelength of 488 nm among the fluorescent proteins of the present invention, a filter having an excitation light between approximately 440 and 460 nm and a fluorescence between approximately 480 and 520 nm can be preferably used. In the case of the fluorescent protein having the excitation maximum wavelength of 548 nm and the fluorescence maximum wavelength of 561 nm among the fluorescent proteins of the present invention, a filter having an excitation light between approximately 530 and 550 nm and a fluorescence between approximately 550 and 600 nm can be preferably used.

[0084] When viable cells are observed over time using a fluorescence microscope, a high sensitive cooled CCD camera is used, since photography is carried out in a short time. In the case of the cooled CCD camera, CCD is cooled to decrease thermal noise, so that a weak fluorescence image can be clearly photographed by exposure in a short time.

(6) Kit of the present invention

[0085] The present invention provides a kit for analyzing the localization of intracellular components and/or analyzing physiologically active substances, which is characterized in that it comprises at least one selected from the fluorescent protein, the fusion fluorescent protein, the DNA, the recombinant vector, or the transformant, which are described in the present specification. The kit of the present invention can be produced from commonly used materials that are known per se, by using common methods.

[0086] Reagents such as the fluorescent protein or the DNA are dissolved in an appropriate solvent, so that the reagents can be prepared in a form suitable for conservation. Water, ethanol, various types of buffer solution, etc. can be used as such a solvent.

[0087] The present invention will be further described in the following examples. However, the present invention is not limited by these examples.

EXAMPLES

Example 1: Isolation of a novel fluorescent protein gene from coral (*Fungia sp.*)

(1) Extraction of total RNA

[0088] A fluorescent protein gene was isolated from coral which emits a fluorescence. 2 individuals of *Fungia sp.* which have red or orange oral disc, were used as a material. *Fungia sp.* was crushed with a hammer. 7.5 ml of "TRIzol" (GIBCO BRL) was added to 4 g (wet weight) of the crushed coral, and the mixture was stirred, followed by centrifugal separation at 1,500 x g for 10 minutes. 1.5 ml of chloroform was added to the obtained supernatant, and the mixture was stirred for 15 seconds, followed by leaving at rest for 3 minutes. Thereafter, the mixture was centrifuged at 7,500 x g for 15 minutes. 3.75 ml of isopropanol was added to the obtained supernatant, and the mixture was stirred for 15 seconds, followed by leaving at rest for 10 minutes. Thereafter, the mixture was centrifuged at 17,000 x g for 10 minutes. The obtained supernatant was discarded. 6 ml of 70% ethanol was added to the residue, and the mixture was centrifuged at 17,000 x g for 10 minutes. The obtained supernatant was discarded, and the precipitate was then dissolved in 200 µl of DEPC water. Total RNA dissolved in the DEPC water was diluted 100 times, and the O.D.260 and O.D. 280 values were measured to determine RNA concentration. 51.6 µg of the total RNA was obtained from the red individual, and 70 µg of the total RNA was obtained from the orange individual.

(2) Synthesis of first strand cDNA

[0089] cDNA (33 µl) was synthesized from 3 µg of the total RNA using a kit for synthesizing first strand cDNA "Ready To Go" (Amersham Pharmacia).

(3) Degenerated PCR

[0090] PCR was carried out using 3 µl of the synthesized first strand cDNA (33 µl) as a template. Primers were designed and produced such that regions similar to the amino acid sequences of known fluorescent proteins were picked up and such regions were converted into nucleotide sequences. The sequences of the used primers are shown below:

5'- GAAGGRTGYGTCAAYGGRCA Y-3' (primer 1) (SEQ ID NO: 9);

5 and

5'- ACVGGDCCATYDGVAAGAAARTT-3' (primer 2) (SEQ ID NO: 10).

10

wherein R represents A or G, Y represents C or T, V represents A, C or G, and D represents A, G or T.

[0091] A PCR reaction solution having the following composition was used:

15

| | |
|------------------------------|------------|
| Template (first strand cDNA) | 3 μ l |
| x10 taq buffer | 5 μ l |
| 2.5 mM dNTPs | 4 μ l |
| 100 μ M primer 1 | 1 μ l |
| 100 μ M primer 2 | 1 μ l |
| Milli-Q | 35 μ l |
| Taq polymerase (5U/ μ l) | 1 μ l |

20

[0092] The following PCR reaction conditions were applied:

25

94°C x 1 minute (PAD)

94°C x 3 seconds (Denaturation)

52°C x 30 seconds (Annealing to template)

72°C x 1 minute (Primer elongation)

30

[0093] A cycle consisting of the above 3 steps was repeated 30 times. The annealing temperature was decreased by 0.3 °C each cycle. This is to say, the temperature was 43°C when 30 cycles were completed.

72°C x 7 minutes (Final elongation)

4°C (Retention)

35

[0094] Using 1 μ l of the amplified product obtained in the first PCR reaction as a template, PCR was carried out again under the same conditions. A 350 bp band of the expected size was cut out and purified by agarose gel electrophoresis.

(4) Subcloning and sequencing

40

[0095] The purified DNA fragment was ligated to a pT7-blue vector (Novagen). *Escherichia coli* (TG1) was transformed therewith and then subjected to blue white selection. Plasmid DNA was purified from white colonies *Escherichia coli*, and the nucleotide sequence of the inserted DNA fragment was determined using a DNA sequencer. Thereafter, the obtained nucleotide sequence was compared with the nucleotide sequences of other fluorescent protein genes, so as to judge whether or not the nucleotide sequence of the DNA fragment was derived from a fluorescent protein.

45

With regard to those that were judged to be a part of the fluorescent protein genes, the full-length gene was cloned by the 5'-RACE method and the 3'-RACE method.

(5) 5'-RACE method

50

[0096] In order to determine a nucleotide sequence on the 5'-side to the DNA fragment obtained by the Degenerated PCR, the 5'-RACE method was applied using 5'-RACE System for Rapid Amplification of cDNA Ends, Version 2.0 (GIBCO BRL). 3 μ g of the total RNA prepared in (1) above was used as a template.

[0097] The following primers were used in the first amplification of dC-tailed cDNA of the orange individual:

55

5'-GGCCACGCGTCGACTAGTACGGGUGGGUGGGUG-3' (SEQ ID NO: 11);

and

5'-GGCTTATATGCGCACTGACTGC-3' (SEQ ID NO: 12)

wherein I represents inosine.

[0098] The following primers were used in the second amplification:

5'-GGCCACGCGTCGACTAGTAC-3' (SEQ ID NO: 13);

and

5'-TATCTCTTCAGGATATTTAGT -3' (SEQ ID NO: 14).

PCR reaction conditions were applied in accordance with protocols attached to the kit.

[0099] The amplified 700 bp band was cut out and purified by agarose gel electrophoresis. The purified DNA fragment was ligated to a pT7-blue vector (Novagen). *Escherichia coli* (TG1) was transformed therewith and then subjected to blue white selection. Plasmid DNA was purified from white colonies *Escherichia coli*, and the nucleotide sequence of the inserted DNA fragment was determined using a DNA sequencer.

[0100] In the same way, the following primers were used in the first amplification of dC-tailed cDNA of the red individual:

5'-GGCCACGCGTCGACTAGTACGGGGIIGGGIIGGGIIG-3' (SEQ ID NO: 15);

and

5'-GGGAAAAGTGCCTTCAATGG-3' (SEQ ID NO: 16)

wherein I represents inosine.

[0101] The following primers were used in the second amplification:

5'- GGCCACGCGTCGACTAGTAC -3' (SEQ ID NO: 17);

and

5'- TCTTCGAACTCAAAC TTTCT -3' (SEQ ID NO: 18).

PCR reaction conditions were applied in accordance with protocols attached to the kit.

[0102] The amplified 500 bp band was cut out and purified by agarose gel electrophoresis. The purified DNA fragment was ligated to a pT7-blue vector (Novagen). *Escherichia coli* (TG1) was transformed therewith and then subjected to blue white selection. Plasmid DNA was purified from white colonies *Escherichia coli*, and the nucleotide sequence of the inserted DNA fragment was determined using a DNA sequencer.

(6) 3'-RACE method

[0103] A nucleotide sequence on the 3'-side to the DNA fragment obtained by the Degenerated PCR was obtained by PCR, using the primer prepared based on the information obtained by determination of the nucleotide sequence in (4) above and an oligo dT primer. 3 µl of the first strand cDNA prepared in (2) above was used as a template.

[0104] The prepared primers were;

For orange individual; 5'- GCAGTCAGTGCGCATATAAGCC -3' (primer3)(SEQ ID NO: 19)

For red individual; 5'- CCATTGAAGGCACTTTTCCC -3'(primer4) (SEQ ID NO: 20)

[0105] A PCR reaction solution having the following composition was used:

| | |
|------------------------------|-------|
| Template (first strand cDNA) | 3 µl |
| x10 taq buffer | 5 µl |
| 2.5 mM dNTPs | 4 µl |
| 20 µM primer 3 or primer 4 | 1 µl |
| 10 µM oligo dT primer | 1 µl |
| Milli-Q | 35 µl |
| Taq polymerase (5U/µl) | 1 µl |

[0106] The following PCR reaction conditions were applied:

94°C x 1 minute (PAD)
 94°C x 30 seconds (Denaturation)
 55°C x 30 seconds (Annealing to template)
 72°C x 1 minute (Primer elongation)

[0107] A cycle consisting of the above 3 steps was repeated 30 times.

72°C x 7 minutes (Final elongation)

4°C (Retention)

[0108] The amplified 850 bp band was cut and purified by agarose gel electrophoresis. The purified DNA fragment was ligated to a pT7-blue vector (Novagen). *Escherichia coli* (TG1) was transformed therewith and then subjected to blue white selection. Plasmid DNA was purified from white colonies *Escherichia coli*, and the nucleotide sequence of the inserted DNA fragment was determined using a DNA sequencer.

(7) Expression of protein in *Escherichia coli*

[0109] From the obtained full-length nucleotide sequence, primers corresponding to the N-terminus and C-terminus of the protein were prepared. Thereafter, PCR was carried out using the first strand cDNA prepared in (2) above as a template. The used primer for both orange and red individuals is primer 5:

5'-CGGGATCCATGAAGATGAAGTACTTTATGGATGG -3'(primer5) (SEQ ID NO: 21)

[0110] A PCR reaction solution having the following composition was used:

| | |
|------------------------------|------|
| Template (first strand cDNA) | 3 µl |
| x10 pyrobest buffer | 5 µl |

(continued)

| | |
|-----------------------------|-------|
| 2.5 mM dNTPs | 4 µl |
| 20 µM primer 5 | 1 µl |
| 20 µM oligo dT primer | 1 µl |
| Milli-Q | 35 µl |
| Pyrobest polymerase (5U/µl) | 1 µl |

[0111] The following PCR reaction conditions were applied:

94°C x 1 minute (PAD)
 94°C x 30 seconds (Denaturation)
 55°C x 30 seconds (Annealing to template)
 72°C x 1 minute (Primer elongation)

[0112] A cycle consisting of the above 3 steps was repeated 30 times.

72°C x 7 minutes (Final elongation)

4°C (Retention)

[0113] An amplified band with a size of approximately 1,000 bp was cut out and purified by agarose gel electrophoresis, and was then subcloned into the BamHI-EcoRI site of a pRSET vector (Invitrogen). Thereafter, it was expressed in *Escherichia coli* (JM109-DE3). However, whatever it may be derived from an orange individual or red individual, the protein expressed in the *Escherichia coli* did not emit fluorescence.

[0114] It was found that proteins translated from the genes cloned herein (a translated protein derived from an orange individual is referred to as Kusabira-Orange, and a translated protein derived from a red individual is referred to as Kusabira-Cyan) lack about 10 amino acids at their N-terminus, when compared with the known fluorescent proteins (FP486 and Azami-Green) (Table 1). Accordingly, 2 types of segments each consisting of about 10 amino acids were added to their N-termini.

Table 1 Comparison of amino acid sequences at N-termini among the fluorescent proteins

Fluorescent proteins derived from orange individuals

↓

Kusabira-Orange-1 (KO-1) *MSVIKPEMKMKYFMDGSVNGHFFTVEGEG...*

-Orange-2 (KO-2) *MALSNKFIGDDMKMKYFMDGSVNGHFFTVEGEG...*

Fluorescent proteins derived from red individuals

Kusabira-Cyan-1 (KCy-1) *MSVIKPEMKMKYFMDGSVNGHFFTVEGEG...*

-Cyan-2 (KCy-2) *MALSNKFIGDDMKMKYFMDGSVNGHFFTVEGEG...*

Azami-Green *MSVIKPEMKIKLCMRGTVNGHNFVIEGEG...*

FP486 *MALSNKFIGDDMKMTYHMDGCVNGHYFTVEGEG...*

[0115] The sequence portions of Kusabira-Orange and Kusabira-Cyan indicated with italicized letters (from the letters pointed with an arrow and downward) have been cloned.

[0116] In one case, 11 amino acids (MALSNKFIGDD) at the N-terminus of FP486 were used, the amino acid se-

quence as a whole of which is similar to those of the above fluorescent proteins. In another case, 7 amino acids (MSVIKPE) at the N-terminus of Azami-Green were used, which had previously been cloned. As a result, it was found that, in both cases where either one of the above amino acid sequences was added, orange fluorescence was emitted from a fluorescent protein derived from an orange individual, and that cyan fluorescence was emitted from a fluorescent protein derived from a red individual. A fluorescent protein produced by adding the 7 amino acids at the N-terminus of Azami-Green to a fluorescent protein derived from an orange individual was defined as Kusabira-Orange-1 (KO-1) (SEQ ID NO: 3), and a fluorescent protein produced by adding the 11 amino acids at the N-terminus of FP486 to a fluorescent protein derived from an orange individual was defined as Kusabira-Orange-2 (KO-2) (SEQ ID NO: 4). A fluorescent protein produced by adding the 7 amino acids at the N-terminus of Azami-Green to a fluorescent protein derived from a red individual was defined as Kusabira-Cyan-1 (KCy-1) (SEQ ID NO: 1), and a fluorescent protein produced by adding the 11 amino acids at the N-terminus of FP486 to a fluorescent protein derived from a red individual was defined as Kusabira-Cyan-2 (KCy-2) (SEQ ID NO: 2).

[0117] The entire nucleotide sequences of KCy-1, KCy-2, KO-1 and KO-2 are shown in SEQ ID NOS: 5 to 8, respectively.

[0118] The above 4 types of proteins were constructed such that His-tag was attached to their N-terminus. Thus, the expressed proteins were purified with Ni-Agarose gel (QIAGEN). Purification was carried out in accordance with protocols attached with the above product. Subsequently, the properties of the purified proteins were analyzed.

(8) Analysis of fluorescence properties

[0119] An absorption spectrum was measured using a 20 μ M fluorescent protein and a 50 mM HEPES solution of pH 7.5. A molar absorption coefficient was calculated from the value of the peak of this spectrum. In the case of each of KCy-1 and KCy-2, the fluorescent protein was diluted with the above buffer solution such that its absorption peak was observed at 455 nm and that its absorption at 400 nm became 0.005. Thereafter, a fluorescence spectrum obtained by excitation at 400 nm and an excitation spectrum obtained by fluorescence at 520 nm were determined. Likewise, ECFP (Clontech) was diluted with the above buffer solution such that its absorption at 400 nm became 0.005, and its fluorescence spectrum was determined. Setting the quantum yield of ECFP at 0.4, the quantum yield of each of KCy-1 and KCy-2 was obtained. In the case of each of KO-1 and KO-2, the fluorescent protein was diluted with the above buffer solution such that its absorption peak was observed at 548 nm and that its absorption at 500 nm became 0.0025. Thereafter, a fluorescence spectrum obtained by excitation at 500 nm and an excitation spectrum obtained by fluorescence at 590 nm were determined. Likewise, DsRed (Clontech) was diluted with the above buffer solution such that its absorption at 500 nm became 0.0025, and its fluorescence spectrum was determined. Setting the quantum yield of DsRed at 0.29, the quantum yield of each of KO-1 and KO-2 was obtained. The results are shown in Table 2 and Figure 1.

Table 2

Comparison of Kusabira-Cyan with ECFP (Clontech)

| | Excitation maximum | Fluorescence maximum | Molar absorption coefficient | Quantum yield | pH stability | Number of amino acids |
|-------|-----------------------|-------------------------|------------------------------------|------------------|------------------------|--------------------------|
| KCy-1 | 455 nm | 488 nm | 38,700 (455 nm) | 0.85 | Stable at pH 5 to 9 | 223 |
| KCy-2 | 455 nm | 488 nm | 27,700 (455 nm) | 0.81 | Stable at pH 5 to 9 | 227 |
| ECFP | 435 nm | 478 nm | 28,750 (435 nm) | 0.40 | pKa = 5.5 | 239 |

Comparison of Kusabira-Orange with DsRed (Clontech)

| | Excitation maximum | Fluorescence maximum | Molar absorption coefficient | Quantum yield | pH stability | Number of amino acids |
|-------|-----------------------|-------------------------|------------------------------------|------------------|--------------|--------------------------|
| KO-1 | 548 nm | 561 nm | 75,900 (548 nm) | 0.44 | pKa < 5.0 | 217 |
| KO-2 | 548 nm | 561 nm | 51,000 (548 nm) | 0.50 | pKa < 5.0 | 221 |
| DsRed | 559 nm | 583 nm | 86,100 (559 nm) | 0.29 | None | 226 |

(9) Difference in expression caused by difference of amino acid sequence at N-terminus

[0120] The above each protein was expressed in *Escherichia coli* (JM109-DE3). 0.1 mM IPTG was used to induce the expression of the protein, and sampling was then carried out thereon at 17 hours, 50 hours, and 74 hours after the expression induction. Thereafter, a change in the fluorescence spectrum was analyzed with a fluorospectrophotometer. The results are shown in Figures 2 and 3. The appearance of fluorescence was different depending on an amino acid sequence added to the N-terminus. In both cases of Kusabira-Orange and Kusabira-Cyan, a protein to which the 7

amino acids at the N-terminus of an Azami-Green (*Galaxea fascicularis* fluorescent protein) were added, emitted fluorescence more rapidly than the other did.

[0121] In the cases of KO-1 and KO-2, green fluorescence was first emitted, and it was then changed to orange fluorescence. Such a maturation process was carried out more rapidly in KO-1 than in KO-2.

(10) Measurement of pH sensitivity

[0122] In the cases of KCy-1 and KCy-2, the fluorescent protein was diluted with the following buffer such that the absorption at 400 nm became 0.005. In the cases of KO-1 and KO-2, the fluorescent protein was diluted with the following buffer such that the absorption at 500 nm became 0.0025. Then, the fluorescence spectrums were measured. The measurement results are shown in Figure 4.

pH 4 and 5: Acetate buffer

pH 6: MES buffer

pH 7: MOPS buffer

pH 8: HEPES buffer

pH 9 and 10: Glycine buffer

pH 11: Phosphate buffer

(Example 2) Production of amino acid-substituted mutant of Kusabira-Orange (KO)

[0123] Kusabira-Orange (KO) is a fluorescent protein which emits orange fluorescence (fluorescence maximum at 561 nm, and excitation maximum at 548 nm). Cysteine at position 64 was substituted by alanine in the amino acid sequence of KO-1, so as to obtain a mutant which emits green fluorescence (fluorescence maximum at 508 nm, and excitation maximum at 496 nm), wherein fluorescent properties were sifted on the short wavelength side when compared with KO (Figure 5). Moreover, glutamic acid at position 211 was substituted by alanine in the amino acid sequence of KO-1, so as to obtain a mutant which emits red fluorescence (fluorescence maximum at 578 nm, and excitation maximum at 563 nm), wherein fluorescent properties were sifted on the long wavelength side when compared with KO (Figure 6).

INDUSTRIAL APPLICABILITY

[0124] The present invention provides novel fluorescent proteins derived from organisms other than a jellyfish. Since the fluorescent proteins of the present invention have desired fluorescent properties and has low pH sensitivity, it is useful in the molecular biological analysis.

SEQUENCE LISTING

5 <110> RIKEN
 <120> Fluorescent proteins
 <130> A21741A
 10 <160> 21

 15 <210> 1
 <211> 223
 <212> PRT
 20 <213> Fungia sp.
 <400> 1
 25 Met Ser Val Ile Lys Pro Glu Met Lys Met Lys Tyr Phe Met Asp Gly
 1 5 10 15
 Ser Val Asn Gly His Glu Phe Thr Val Glu Gly Glu Gly Thr Gly Lys
 30 20 25 30
 Pro Tyr Glu Gly Lys His Lys Ile Thr Leu Asp Val Thr Lys Gly Gly
 35 35 40 45
 Pro Leu Pro Phe Ala Phe Asp Leu Leu Ser Thr Val Phe Ser Tyr Gly
 50 55 60
 40 Asn Arg Cys Leu Thr Lys Tyr Pro Asp Asp Ile Pro Asp Tyr Phe Lys
 65 70 75 80
 45 Gln Cys Phe Pro Gly Gly Tyr Ser Trp Glu Arg Lys Phe Glu Phe Glu
 85 90 95
 Asp Gly Gly Leu Ala Ile Ala Lys Ala Glu Ile Ser Leu Lys Gly Asn
 50 100 105 110
 Cys Phe Glu His Lys Ser Thr Ile Glu Gly Thr Phe Pro Asp Ser Ser
 55 115 120 125

5 Pro Ile Ala Gln Asn Lys Thr Leu Gly Trp Glu Pro Ser Thr Glu Lys
130 135 140

Met Thr Val Arg Asp Gly Ser Met Lys Gly Asp Asp Ala Ala Tyr Leu
10 145 150 155 160

Lys Leu Val Gly Gly Gly Asn His Lys Cys Tyr Phe Thr Thr Thr Tyr
15 165 170 175

Thr Ala Lys Lys Lys Ile Pro Asn Leu Pro Gln Ser His Phe Ile Gly
180 185 190

20 His Arg Ile Ser Ser Val Val Asn Gly Thr Lys Ile Gly Val Met Glu
195 200 205

Asp Ala Ile Ala His Leu Tyr Pro Phe Asn Gly Val Pro Cys Gln
25 210 215 220

30 <210> 2
<211> 227
<212> PRT

35 <213> Fungia sp.
<400> 2

40 Met Ala Leu Ser Asn Lys Phe Ile Gly Asp Asp Met Lys Met Lys Tyr
1 5 10 15

Phe Met Asp Gly Ser Val Asn Gly His Glu Phe Thr Val Glu Gly Glu
45 20 25 30

Gly Thr Gly Lys Pro Tyr Glu Gly Lys His Lys Ile Thr Leu Asp Val
50 35 40 45

Thr Lys Gly Gly Pro Leu Pro Phe Ala Phe Asp Leu Leu Ser Thr Val
50 55 60

55 Phe Ser Tyr Gly Asn Arg Cys Leu Thr Lys Tyr Pro Asp Asp Ile Pro

EP 1 464 701 A1

| | | | | |
|----|-----|-----|-----|-----|
| 5 | 65 | 70 | 75 | 80 |
| | Asp | Tyr | Phe | Lys |
| | Gln | Cys | Phe | Pro |
| | Gly | Gly | Tyr | Ser |
| | Trp | Glu | Arg | Lys |
| | 85 | 90 | 95 | |
| 10 | Phe | Glu | Phe | Glu |
| | Asp | Gly | Gly | Leu |
| | Ala | Ile | Ala | Lys |
| | Ala | Glu | Ile | Ser |
| | 100 | 105 | 110 | |
| | Leu | Lys | Gly | Asn |
| | Cys | Phe | Glu | His |
| | Lys | Ser | Thr | Ile |
| | Glu | Gly | Thr | Phe |
| 15 | 115 | 120 | 125 | |
| | Pro | Asp | Ser | Ser |
| | Pro | Ile | Ala | Gln |
| | Asn | Lys | Thr | Leu |
| | Gly | Trp | Glu | Pro |
| 20 | 130 | 135 | 140 | |
| | Ser | Thr | Glu | Lys |
| | Met | Thr | Val | Arg |
| | Asp | Gly | Ser | Met |
| | Lys | Gly | Asp | Asp |
| | 145 | 150 | 155 | 160 |
| 25 | Ala | Ala | Tyr | Leu |
| | Lys | Leu | Val | Gly |
| | Gly | Gly | Gly | Asn |
| | His | Lys | Cys | Tyr |
| | Phe | 165 | 170 | 175 |
| | Thr | Thr | Thr | Tyr |
| | Thr | Ala | Lys | Lys |
| | Lys | Lys | Ile | Pro |
| | Asn | Leu | Pro | Gln |
| | Ser | 180 | 185 | 190 |
| | His | Phe | Ile | Gly |
| | His | Arg | Ile | Ser |
| | Ser | Val | Val | Asn |
| | Gly | Thr | Lys | Ile |
| 35 | 195 | 200 | 205 | |
| | Gly | Val | Met | Glu |
| | Asp | Ala | Ile | Ala |
| | His | Leu | Tyr | Pro |
| | Phe | Asn | Gly | Val |
| | 210 | 215 | 220 | |
| 40 | Pro | Cys | Gln | |
| | 225 | | | |

45

<210> 3

50

<211> 217

<212> PRT

<213> Fungia sp.

55

<400> 3

EP 1 464 701 A1

| | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 5 | Met | Ser | Val | Ile | Lys | Pro | Glu | Met | Lys | Met | Lys | Tyr | Phe | Met | Asp | Gly |
| | 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| | Ser | Val | Asn | Gly | His | Glu | Phe | Thr | Val | Glu | Gly | Glu | Gly | Thr | Gly | Lys |
| 10 | | | | 20 | | | | | 25 | | | | | | 30 | |
| | Pro | Tyr | Glu | Gly | His | Gln | Glu | Met | Thr | Leu | Arg | Val | Thr | Met | Ala | Lys |
| | | | | 35 | | | | | 40 | | | | | | 45 | |
| 15 | Gly | Gly | Pro | Met | Pro | Phe | Ser | Phe | Asp | Leu | Val | Ser | His | Thr | Phe | Cys |
| | | 50 | | | | | 55 | | | | | 60 | | | | |
| 20 | Tyr | Gly | His | Arg | Pro | Phe | Thr | Lys | Tyr | Pro | Glu | Glu | Ile | Pro | Asp | Tyr |
| | | 65 | | | | 70 | | | | | 75 | | | | 80 | |
| | Phe | Lys | Gln | Ala | Phe | Pro | Glu | Gly | Leu | Ser | Trp | Glu | Arg | Ser | Leu | Gln |
| 25 | | | | | 85 | | | | | 90 | | | | | 95 | |
| | Phe | Glu | Asp | Gly | Gly | Phe | Ala | Ala | Val | Ser | Ala | His | Ile | Ser | Leu | Arg |
| 30 | | | | 100 | | | | | 105 | | | | | | 110 | |
| | Gly | Asn | Cys | Phe | Glu | His | Lys | Ser | Lys | Phe | Val | Gly | Val | Asn | Phe | Pro |
| | | | 115 | | | | | 120 | | | | | 125 | | | |
| 35 | Ala | Asp | Gly | Pro | Val | Met | Gln | Asn | Gln | Ser | Ser | Asp | Trp | Glu | Pro | Ser |
| | | | 130 | | | | 135 | | | | | | 140 | | | |
| 40 | Thr | Glu | Lys | Ile | Thr | Thr | Cys | Asp | Gly | Val | Leu | Lys | Gly | Asp | Val | Thr |
| | | 145 | | | | 150 | | | | | 155 | | | | 160 | |
| | Met | Phe | Leu | Lys | Leu | Ala | Gly | Gly | Gly | Asn | His | Lys | Cys | Gln | Phe | Lys |
| 45 | | | | 165 | | | | | 170 | | | | | | 175 | |
| | Thr | Thr | Tyr | Lys | Ala | Ala | Lys | Lys | Ile | Leu | Lys | Met | Pro | Gln | Ser | His |
| 50 | | | | 180 | | | | | 185 | | | | | | 190 | |
| | Phe | Ile | Gly | His | Arg | Leu | Val | Arg | Lys | Thr | Glu | Gly | Asn | Ile | Thr | Glu |
| | | | 195 | | | | | 200 | | | | | 205 | | | |
| 55 | Leu | Val | Glu | Asp | Ala | Val | Ala | His | Cys | | | | | | | |

210

215

5

<210> 4

10

<211> 221

<212> PRT

<213> Fungia sp.

15

<400> 4

Met Ala Leu Ser Asn Lys Phe Ile Gly Asp Asp Met Lys Met Lys Tyr

20

1 5 10 15

Phe Met Asp Gly Ser Val Asn Gly His Glu Phe Thr Val Glu Gly Glu

20 25 30

25

Gly Thr Gly Lys Pro Tyr Glu Gly His Gln Glu Met Thr Leu Arg Val

35 40 45

30

Thr Met Ala Lys Gly Gly Pro Met Pro Phe Ser Phe Asp Leu Val Ser

50 55 60

His Thr Phe Cys Tyr Gly His Arg Pro Phe Thr Lys Tyr Pro Glu Glu

35

65 70 75 80

Ile Pro Asp Tyr Phe Lys Gln Ala Phe Pro Glu Gly Leu Ser Trp Glu

85 90 95

40

Arg Ser Leu Gln Phe Glu Asp Gly Gly Phe Ala Ala Val Ser Ala His

100 105 110

45

Ile Ser Leu Arg Gly Asn Cys Phe Glu His Lys Ser Lys Phe Val Gly

115 120 125

50

Val Asn Phe Pro Ala Asp Gly Pro Val Met Gln Asn Gln Ser Ser Asp

130 135 140

Trp Glu Pro Ser Thr Glu Lys Ile Thr Thr Cys Asp Gly Val Leu Lys

55

145 150 155 160

EP 1 464 701 A1

Gly Asp Val Thr Met Phe Leu Lys Leu Ala Gly Gly Gly Asn His Lys
 5 165 170 175
 Cys Gln Phe Lys Thr Thr Tyr Lys Ala Ala Lys Lys Ile Leu Lys Met
 10 180 185 190
 Pro Gln Ser His Phe Ile Gly His Arg Leu Val Arg Lys Thr Glu Gly
 195 200 205
 15 Asn Ile Thr Glu Leu Val Glu Asp Ala Val Ala His Cys
 210 215 220
 20
 <210> 5
 <211> 672
 25 <212> DNA
 <213> Fungia sp.
 <220>
 30 <221> CDS
 <222> (1).. (669)
 35 <400> 5
 atg agt gtg att aaa cca gag atg aag atg aag tac ttc atg gac gga 48
 40 Met Ser Val Ile Lys Pro Glu Met Lys Met Lys Tyr Phe Met Asp Gly
 1 5 10 15
 tcc gtc aat ggg cat gag ttc aca gtt gaa ggt gaa ggc aca ggc aaa 96
 45 Ser Val Asn Gly His Glu Phe Thr Val Glu Gly Glu Gly Thr Gly Lys
 20 25 30
 cct tac gag gga aag cac aaa ata aca ctt gac gtc acc aag ggt ggg 144
 50 Pro Tyr Glu Gly Lys His Lys Ile Thr Leu Asp Val Thr Lys Gly Gly
 35 40 45
 55 cca ctg cct ttt gcg ttt gac ttg ttg tct aca gtg ttc tct tat ggc 192

EP 1 464 701 A1

| | | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | Pro | Leu | Pro | Phe | Ala | Phe | Asp | Leu | Leu | Ser | Thr | Val | Phe | Ser | Tyr | Gly | |
| 5 | | | | | | | | | | | | | | | | | |
| | | | 50 | | | | | 55 | | | | | 60 | | | | |
| | aac | aga | tgc | ctt | act | aaa | tat | cct | gac | gat | atc | ccc | gac | tat | ttc | aaa | 240 |
| | Asn | Arg | Cys | Leu | Thr | Lys | Tyr | Pro | Asp | Asp | Ile | Pro | Asp | Tyr | Phe | Lys | |
| 10 | | | | | | | | | | | | | | | | | |
| | | | 65 | | | | | 70 | | | | | 75 | | | 80 | |
| | caa | tgc | ttt | cct | gga | ggc | tat | tca | tgg | gaa | aga | aag | ttt | gag | ttc | gaa | 288 |
| | Gln | Cys | Phe | Pro | Gly | Gly | Tyr | Ser | Trp | Glu | Arg | Lys | Phe | Glu | Phe | Glu | |
| 15 | | | | | | | | | | | | | | | | | |
| | | | | | | 85 | | | | | 90 | | | | 95 | | |
| | gat | ggc | ggg | ttg | gct | ata | gcc | aaa | gcg | gaa | ata | agc | ctt | aaa | gga | aac | 336 |
| 20 | | | | | | | | | | | | | | | | | |
| | Asp | Gly | Gly | Leu | Ala | Ile | Ala | Lys | Ala | Glu | Ile | Ser | Leu | Lys | Gly | Asn | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | 100 | | | | | 105 | | | | 110 | | |
| 25 | | | | | | | | | | | | | | | | | |
| | tgc | ttc | gaa | cac | aaa | tcc | acc | att | gaa | ggc | act | ttt | ccc | gat | agc | agt | 384 |
| | Cys | Phe | Glu | His | Lys | Ser | Thr | Ile | Glu | Gly | Thr | Phe | Pro | Asp | Ser | Ser | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | 115 | | | | | 120 | | | | 125 | | |
| 30 | | | | | | | | | | | | | | | | | |
| | cct | att | gcg | caa | aac | aag | acg | cta | gga | tgg | gaa | cca | tcc | acc | gag | aag | 432 |
| | Pro | Ile | Ala | Gln | Asn | Lys | Thr | Leu | Gly | Trp | Glu | Pro | Ser | Thr | Glu | Lys | |
| | | | | | | | | | | | | | | | | | |
| 35 | | | | | | | | | | | | | | | | | |
| | | | | | | 130 | | | | | 135 | | | | 140 | | |
| | atg | acc | gtc | cgc | gac | gga | tca | atg | aag | ggt | gat | gat | gcg | gcc | tac | ctc | 480 |
| | Met | Thr | Val | Arg | Asp | Gly | Ser | Met | Lys | Gly | Asp | Asp | Ala | Ala | Tyr | Leu | |
| 40 | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | 145 | | | | | 150 | | | | 155 | 160 | |
| | aaa | ttg | gtg | gga | ggc | ggc | aat | cac | aaa | tgc | tac | ttt | aca | act | acc | tac | 528 |
| | Lys | Leu | Val | Gly | Gly | Gly | Asn | His | Lys | Cys | Tyr | Phe | Thr | Thr | Thr | Tyr | |
| 45 | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | 165 | | | | | 170 | | | | 175 | | |
| | aca | gcg | aag | aaa | aag | att | cct | aac | ctg | cca | caa | agc | cat | ttc | att | ggg | 576 |
| 50 | | | | | | | | | | | | | | | | | |
| | Thr | Ala | Lys | Lys | Lys | Ile | Pro | Asn | Leu | Pro | Gln | Ser | His | Phe | Ile | Gly | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| | | | | | | 180 | | | | | 185 | | | | 190 | | |
| 55 | | | | | | | | | | | | | | | | | |
| | cat | cgc | atc | tcc | agt | gtc | gtc | aat | ggc | act | aaa | att | gga | gtg | atg | gaa | 624 |

His Arg Ile Ser Ser Val Val Asn Gly Thr Lys Ile Gly Val Met Glu

5

195

200

205

gat gca att gct cat ctt tac cct ttt aat ggc gtg cca tgc cag tga 672

Asp Ala Ile Ala His Leu Tyr Pro Phe Asn Gly Val Pro Cys Gln

10

210

215

220

15

<210> 6

<211> 684

<212> DNA

20

<213> Fungia sp.

<220>

25

<221> CDS

<222> (1).. (681)

<400> 6

30

atg gcc ctg agc aac aag ttc atc ggg gac gac atg aag atg aag tac 48

Met Ala Leu Ser Asn Lys Phe Ile Gly Asp Asp Met Lys Met Lys Tyr

35

1

5

10

15

ttc atg gac gga tcc gtc aat ggg cat gag ttc aca gtt gaa ggt gaa 96

Phe Met Asp Gly Ser Val Asn Gly His Glu Phe Thr Val Glu Gly Glu

40

20

25

30

ggc aca ggc aaa cct tac gag gga aag cac aaa ata aca ctt gac gtc 144

45

Gly Thr Gly Lys Pro Tyr Glu Gly Lys His Lys Ile Thr Leu Asp Val

35

40

45

acc aag ggt ggg cca ctg cct ttt gcg ttt gac ttg ttg tct aca gtg 192

50

Thr Lys Gly Gly Pro Leu Pro Phe Ala Phe Asp Leu Leu Ser Thr Val

50

55

60

55

ttc tct tat ggc aac aga tgc ctt act aaa tat cct gac gat atc ccc 240



55

| | | | |
|---|-----|-----|-----|
| 210 | 215 | 220 | |
| cca tgc cag tga | | | 684 |
| Pro Cys Gln | | | |
| 225 | | | |
| <210> 7 | | | |
| <211> 654 | | | |
| <212> DNA | | | |
| <213> Fungia sp. | | | |
| <220> | | | |
| <221> CDS | | | |
| <222> (1).. (651) | | | |
| <400> 7 | | | |
| atg agt gtg att aaa cca gag atg aag atg aag tac ttc atg gac gga | | | 48 |
| Met Ser Val Ile Lys Pro Glu Met Lys Met Lys Tyr Phe Met Asp Gly | | | |
| 1 5 10 15 | | | |
| tcc gtc aat ggg cat gag ttc aca gtt gaa ggt gaa ggc aca ggc aaa | | | 96 |
| Ser Val Asn Gly His Glu Phe Thr Val Glu Gly Glu Gly Thr Gly Lys | | | |
| 20 25 30 | | | |
| cct tac gag gga cat caa gag atg aca cta cgc gtc aca atg gcc aag | | | 144 |
| Pro Tyr Glu Gly His Gln Glu Met Thr Leu Arg Val Thr Met Ala Lys | | | |
| 35 40 45 | | | |
| ggc ggg cca atg cct ttc tcg ttt gac tta gtg tca cac acg ttc tgt | | | 192 |
| Gly Gly Pro Met Pro Phe Ser Phe Asp Leu Val Ser His Thr Phe Cys | | | |
| 50 55 60 | | | |
| tac ggc cac aga cct ttt act aaa tat cca gaa gag ata cca gac tat | | | 240 |

EP 1 464 701 A1

| | | |
|----|--|-----|
| | Tyr Gly His Arg Pro Phe Thr Lys Tyr Pro Glu Glu Ile Pro Asp Tyr | |
| 5 | 65 70 75 80 | |
| | ttc aaa caa gca ttt cct gaa ggc ctg tca tgg gaa agg tcg ttg cag | 288 |
| 10 | Phe Lys Gln Ala Phe Pro Glu Gly Leu Ser Trp Glu Arg Ser Leu Gln | |
| | 85 90 95 | |
| | ttc gaa gat ggt ggg ttt gct gca gtc agt gcg cat ata agc ctt aga | 336 |
| 15 | Phe Glu Asp Gly Gly Phe Ala Ala Val Ser Ala His Ile Ser Leu Arg | |
| | 100 105 110 | |
| 20 | gga aac tgc ttc gag cac aaa tcc aaa ttt gtt ggg gtt aac ttt cct | 384 |
| | Gly Asn Cys Phe Glu His Lys Ser Lys Phe Val Gly Val Asn Phe Pro | |
| | 115 120 125 | |
| 25 | gcc gat ggt cct gtg atg caa aac caa agt tct gat tgg gag cca tca | 432 |
| | Ala Asp Gly Pro Val Met Gln Asn Gln Ser Ser Asp Trp Glu Pro Ser | |
| 30 | 130 135 140 | |
| | acc gag aaa att act acc tgc gac gga gtt ctg aag ggt gat gtt acg | 480 |
| | Thr Glu Lys Ile Thr Thr Cys Asp Gly Val Leu Lys Gly Asp Val Thr | |
| 35 | 145 150 155 160 | |
| | atg ttc cta aag ctt gcg gga ggc ggc aat cac aaa tgc caa ttc aag | 528 |
| 40 | Met Phe Leu Lys Leu Ala Gly Gly Gly Asn His Lys Cys Gln Phe Lys | |
| | 165 170 175 | |
| | act act tac aag gcg gca aaa aag att ctt aaa atg cca caa agc cat | 576 |
| 45 | Thr Thr Tyr Lys Ala Ala Lys Lys Ile Leu Lys Met Pro Gln Ser His | |
| | 180 185 190 | |
| 50 | ttc atc ggg cat cgc ctc gtc agg aaa acc gaa ggc aac att act gag | 624 |
| | Phe Ile Gly His Arg Leu Val Arg Lys Thr Glu Gly Asn Ile Thr Glu | |
| | 195 200 205 | |
| 55 | ctg gta gaa gat gca gta gct cat tgc tga | 654 |

Leu Val Glu Asp Ala Val Ala His Cys

5

210

215

<210> 8

10

<211> 666

<212> DNA

15

<213> Fungia sp.

<220>

<221> CDS

20

<222> (1).. (663)

<400> 8

25

atg gcc ctg agc aac aag ttc atc ggg gac gac atg aag atg aag tac 48

Met Ala Leu Ser Asn Lys Phe Ile Gly Asp Asp Met Lys Met Lys Tyr

1

5

10

15

30

ttc atg gac gga tcc gtc aat ggg cat gag ttc aca gtt gaa ggt gaa 96

Phe Met Asp Gly Ser Val Asn Gly His Glu Phe Thr Val Glu Gly Glu

35

20

25

30

ggc aca ggc aaa cct tac gag gga cat caa gag atg aca cta cgc gtc 144

Gly Thr Gly Lys Pro Tyr Glu Gly His Gln Glu Met Thr Leu Arg Val

40

35

40

45

aca atg gcc aag ggc ggg cca atg cct ttc tcg ttt gac tta gtg tca 192

Thr Met Ala Lys Gly Gly Pro Met Pro Phe Ser Phe Asp Leu Val Ser

45

50

55

60

cac acg ttc tgt tac ggc cac aga cct ttt act aaa tat cca gaa gag 240

His Thr Phe Cys Tyr Gly His Arg Pro Phe Thr Lys Tyr Pro Glu Glu

50

65

70

75

80

55

ata cca gac tat ttc aaa caa gca ttt cct gaa ggc ctg tca tgg gaa 288

EP 1 464 701 A1

| | | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | Ile | Pro | Asp | Tyr | Phe | Lys | Gln | Ala | Phe | Pro | Glu | Gly | Leu | Ser | Trp | Glu | |
| 5 | | | | | | 85 | | | | 90 | | | | | 95 | | |
| | agg | tcg | ttg | cag | ttc | gaa | gat | ggt | ggg | ttt | gct | gca | gtc | agt | gcg | cat | 336 |
| 10 | Arg | Ser | Leu | Gln | Phe | Glu | Asp | Gly | Gly | Phe | Ala | Ala | Val | Ser | Ala | His | |
| | | | | | | 100 | | | | 105 | | | | | 110 | | |
| | ata | agc | ctt | aga | gga | aac | tgc | ttc | gag | cac | aaa | tcc | aaa | ttt | gtt | ggg | 384 |
| 15 | Ile | Ser | Leu | Arg | Gly | Asn | Cys | Phe | Glu | His | Lys | Ser | Lys | Phe | Val | Gly | |
| | | | | | | 115 | | | | 120 | | | | | 125 | | |
| 20 | gtt | aac | ttt | cct | gcc | gat | ggt | cct | gtg | atg | caa | aac | caa | agt | tct | gat | 432 |
| | Val | Asn | Phe | Pro | Ala | Asp | Gly | Pro | Val | Met | Gln | Asn | Gln | Ser | Ser | Asp | |
| | | | | | | 130 | | | | 135 | | | | | 140 | | |
| 25 | tgg | gag | cca | tca | acc | gag | aaa | att | act | acc | tgc | gac | gga | gtt | ctg | aag | 480 |
| | Trp | Glu | Pro | Ser | Thr | Glu | Lys | Ile | Thr | Thr | Cys | Asp | Gly | Val | Leu | Lys | |
| 30 | 145 | | | | | 150 | | | | 155 | | | | | 160 | | |
| | ggt | gat | gtt | acg | atg | ttc | cta | aag | ctt | gcg | gga | ggc | ggc | aat | cac | aaa | 528 |
| 35 | Gly | Asp | Val | Thr | Met | Phe | Leu | Lys | Leu | Ala | Gly | Gly | Gly | Asn | His | Lys | |
| | | | | | | 165 | | | | 170 | | | | | 175 | | |
| 40 | tgc | caa | ttc | aag | act | act | tac | aag | gcg | gca | aaa | aag | att | ctt | aaa | atg | 576 |
| | Cys | Gln | Phe | Lys | Thr | Thr | Tyr | Lys | Ala | Ala | Lys | Lys | Ile | Leu | Lys | Met | |
| | | | | | | 180 | | | | 185 | | | | | 190 | | |
| 45 | cca | caa | agc | cat | ttc | atc | ggg | cat | cgc | ctc | gtc | agg | aaa | acc | gaa | ggc | 624 |
| | Pro | Gln | Ser | His | Phe | Ile | Gly | His | Arg | Leu | Val | Arg | Lys | Thr | Glu | Gly | |
| | | | | | | 195 | | | | 200 | | | | | 205 | | |
| 50 | aac | att | act | gag | ctg | gta | gaa | gat | gca | gta | gct | cat | tgc | tga | 666 | | |
| | Asn | Ile | Thr | Glu | Leu | Val | Glu | Asp | Ala | Val | Ala | His | Cys | | | | |
| 55 | | | | | | 210 | | | | 215 | | | | | 220 | | |

5 <210> 9
 <211> 21
 <212> DNA
 10 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA
 15 <400> 9
 gaaggrtgyg tcaayggrca y 21
 20
 <210> 10
 <211> 23
 25 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <223> Description of Artificial Sequence: Synthetic DNA
 35 <400> 10
 acvggdccat ydgvaagaaa rtt 23
 40
 <210> 11
 <211> 36
 45 <212> DNA
 <213> Artificial Sequence
 <220>
 50 <223> Description of Artificial Sequence: Synthetic DNA
 <400> 11
 55 ggccacgcgt cgactagtac gggiigggii gggiig 36

<210> 12

5

<211> 22

<212> DNA

10

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

15

<400> 12

ggcttatatg cgcactgact gc

22

20

<210> 13

<211> 20

25

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence: Synthetic DNA

35

<400> 13

ggccacgcgt cgactagtag

20

40

<210> 14

<211> 21

45

<212> DNA

<213> Artificial Sequence

<220>

50

<223> Description of Artificial Sequence: Synthetic DNA

<400> 14

55

tatctcttca ggatatttag t

21

<210> 15

5

<211> 36

<212> DNA

10

<213> Artificial Sequence

<220>

15

<223> Description of Artificial Sequence: Synthetic DNA

<400> 15

ggccacgcgt cgactagtac gggiigggi gggiig 36

20

<210> 16

<211> 20

25

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 16

35

gggaaaagtg ccttcaatgg 20

40

<210> 17

<211> 20

45

<212> DNA

<213> Artificial Sequence

<220>

50

<223> Description of Artificial Sequence: Synthetic DNA

<400> 17

55

ggccacgcgt cgactagtac 20

<210> 18

5

<211> 20

<212> DNA

10

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

15

<400> 18

tcttcgaact caaactttct

20

20

<210> 19

<211> 22

25

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence: Synthetic DNA

35

<400> 19

gcagtcagtg cgcatataag cc

22

40

<210> 20

<211> 20

45

<212> DNA

<213> Artificial Sequence

<220>

50

<223> Description of Artificial Sequence: Synthetic DNA

<400> 20

55

ccattgaagg cacttttccc

20

<210> 21

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 21

cgggatccat gaagatgaag tactttatgg atgg 34

Claims

1. A fluorescent protein derived from *Fungia sp.*, which has the following properties:

- (1) the excitation maximum wavelength is 455 nm, and the fluorescence maximum wavelength is 488 nm;
- (2) the molar absorption coefficient at 455 nm is 38700 or 27700;
- (3) the quantum yield is 0.85 or 0.81; and
- (4) the pH sensitivity of the fluorescent property is stable at pH 5 to 9.

2. A fluorescent protein derived from *Fungia sp.*, which has the following properties:

- (1) the excitation maximum wavelength is 548 nm, and the fluorescence maximum wavelength is 561 nm;
- (2) the molar absorption coefficient at 548 nm is 75900 or 51000;
- (3) the quantum yield is 0.44 or 0.50; and
- (4) the pH sensitivity of the fluorescent property is pKa < 5.0.

3. A fluorescent protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1 or 2, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2.

4. A fluorescent protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 3 or 4, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4.

5. A protein having an amino acid sequence wherein cysteine at position 64 is substituted with alanine with respect to the amino acid sequence shown in SEQ ID NO: 3, or an amino acid sequence wherein glutamic acid at position 211 is substituted with alanine with respect to the amino acid sequence shown in SEQ ID NO: 3.

6. A DNA which encodes the fluorescent protein of any one of claims 1 to 5.

7. A DNA which encodes the protein of the following (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1 or 2, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 1 or 2.

8. A DNA which encodes the protein of (a) or (b):

- (a) a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4; or
- (b) a protein having an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 3 or 4, and having fluorescent properties which are equivalent to those of a protein having an amino acid sequence shown in SEQ ID NO: 3 or 4.

9. A DNA of the following (a) or (b):

- (a) DNA having a nucleotide sequence shown in SEQ ID NO: 5 or 6; or
- (b) DNA having a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucleotides with respect to the nucleotide sequence shown in SEQ ID NO: 5 or 6, and encoding a protein having fluorescent properties which are equivalent to those of a protein encoded by the nucleotide sequence shown in SEQ ID NO: 5 or 6.

10. A DNA of the following (a) or (b):

- (a) DNA having a nucleotide sequence shown in SEQ ID NO: 7 or 8; or
- (b) DNA having a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucleotides with respect to the nucleotide sequence shown in SEQ ID NO: 7 or 8, and encoding a protein having fluorescent properties which are equivalent to those of a protein encoded by the nucleotide sequence shown in SEQ ID NO: 7 or 8.

11. A recombinant vector having the DNA of any one of claims 6 to 10.

12. A transformant having the DNA of any one of claims 6 to 10 or the recombinant vector of claim 11.

13. A fusion fluorescent protein consisting of the fluorescent protein of any one of claims 1 to 5 and another protein.

14. The fusion protein of claim 13 wherein said another protein is one that localizes in the cell.

15. The fusion protein of claim 13 or 14 wherein said another protein is one specific to an intracellular organella.

16. A method for analyzing the localization or dynamics of a protein in cells, **characterized in that** the fusion protein of any one of claims 13 to 15 is allowed to be expressed in cells.

17. A fluorescent reagent kit which comprises the fluorescent protein of any one of claims 1 to 5, the DNA of any one of claims 6 to 10, the recombinant vector of claim 11, the transformant of claim 12, or the fusion protein of any one of claims 13 to 15.

Fig.1







Fig.2



2-1



2-2



2-3

Fig.3




Fig. 4




Fig.5






Fig.6



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/13363

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl⁷ C12N15/09, C07K14/435, C07K19/00, G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl⁷ C12N15/09, C07K14/435, C07K19/00, G01N33/68

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 REGISTRY (STN), CA (STN), MEDLINE (STN), WPI (DIALOG), BIOSIS (DIALOG),
 JICST FILE (JOIS), GenBank/EMBL/DBJ/GeneSeq, SwissProt/PIR/GeneSeq

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| Y | MATZ, MV. et al., Fluorescent proteins from nonbioluminescent Anthozoa species. Nat. Biotechnol. (1999), Vol.17, No.10, pages 969 to 973 | 1-17 |
| Y | LUKYANOV, KA. et al., Natural animal coloration can be determined by a nonfluorescent green fluorescent protein homolog. J.Biol.Chem. (2000), Vol.275, No.34, pages 25879 to 25882 | 1-17 |
| Y | Takahiro HOSAKA, "Sango kara Hakken sareta Arata na Keikosei Tanpakushitsu", Kagaku to Kogyo, (2000), Vol.53, No.5, page 612 | 1-17 |
| Y | WO 01/27150 A2 (CLONTECH LAB INC.), 19 April, 2001 (19.04.01), Full text & AU 200110867 A | 1-17 |

☒ Further documents are listed in the continuation of Box C.☐ See patent family annex.

| | |
|---|--|
| * Special categories of cited documents: | "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| "A" document defining the general state of the art which is not considered to be of particular relevance | "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| "E" earlier document but published on or after the international filing date | "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | "&" document member of the same patent family |
| "O" document referring to an oral disclosure, use, exhibition or other means | |
| "P" document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search
03 March, 2003 (03.03.03)Date of mailing of the international search report
25 March, 2003 (25.03.03)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/13363

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|-----------------------|
| Y | WO 00/34320 A2 (CLONTECH LAB INC.), 15 June, 2000 (15.06.00), Full text (Family: none) | 1-17 |
| Y | WO 00/34526 A1 (CLONTECH LAB INC.), 15 June, 2000 (15.06.00), Full text & EP 1135532 A1 & JP 2002-531146 A1 | 1-17 |
| Y | FRADKOV, AF. et al., Novel fluorescent protein from Discosoma coral and its mutants possesses a unique far-red fluorescence. FEBS Lett. (2000), Vol.479, No.3, pages 127 to 130 | 1-17 |
| Y | HEIKAL, AA. et al., Molecular spectroscopy and dynamics of intrinsically fluorescent proteins: coral red (dsRed) and yellow (Citrine). Proc.Natl.Acad.Sci.USA. (2000), Vol.97, No.22, pages 11996 to 12001 | 1-17 |
| Y | Oz Reef Press. Resident of the Month. [online]. Oz Reef Marine Park. (June. 1998), [retrieved on 2003-03-03]. Retrieved from the Internet. <URL: http://ozreef.org/press/1998/june.html#RESIDENT>, <URL:http://web.archive.org/web/20000524121632/ http://ozreef.org/press/1998/june.html> | 1-17 |
| Y | Oz Reef Press. Resident of the Month. [online]. Oz Reef Marine Park. (May. 1997), [retrieved on 2003-03-03]. Retrieved from the Internet. <URL: http://ozreef.org/press/1997/may.html#RESIDENT>, <URL:http://web.archive.org/web/20000602051932/ http://ozreef.org/press/1997/may.html> | 1-17 |

Form PCT/ISA/210 (continuation of second sheet) (July 1998)